

ACOG RECOMMENDATIONS AND THE LAMAZE SIX HEALTHY BIRTH PRACTICES

Lamaze Six Healthy Birth Practices	Recommendations of the American College of Obstetricians and Gynecologists (ACOG)
<p>1 Let labor begin on its own.</p>	<p>Before the publication of the ARRIVE Trial (A Randomized Trial of Induction versus Expectant Management)¹ in the <i>New England Journal of Medicine</i> in August of 2018, ACOG discouraged elective induction before 41 -42 weeks of gestation. In response to the ARRIVE Trial, ACOG issued a practice advisory² stating that, “it is reasonable for obstetricians and health care facilities to offer elective induction of labor to low-risk nulliparous [first-time] women at 39 weeks gestation,” provided that</p> <ul style="list-style-type: none"> • The due date has been confirmed by an ultrasound early in the pregnancy,³ • Thorough discussion has taken place between the provider and pregnant woman² and • Ample time is provided for labor to progress. ACOG recommends “allowing longer durations for latent labor (up to 24 hours or longer) and requiring that oxytocin be administered for at least 12-18 hours after membrane rupture before deeming the induction a failure.”²
<p>2 Walk, move around, and change positions throughout labor.</p>	<p>“Frequent position changes during labor to enhance maternal comfort and promote optimal fetal positioning can be supported as long as adopted positions allow appropriate maternal and fetal monitoring and treatments and are not contraindicated by maternal medical or obstetric complications.”⁴</p>
<p>3 Bring a loved one, friend, or doula for continuous support.</p>	<p>“Published data indicate that one of the most effective tools to improve labor and delivery outcomes is the continuous presence of support personnel, such as a doula.”⁵</p> <ul style="list-style-type: none"> • “Evidence suggests that, in addition to regular nursing care, continuous one-to-one emotional support is associated with improved outcomes for women in labor.”³
<p>4 Avoid interventions that are not medically necessary.</p>	<ul style="list-style-type: none"> • “...for women with normally progressing labor and no evidence of fetal compromise, routine amniotomy need not be undertaken unless required to facilitate monitoring.”⁴ • “To facilitate the option of intermittent auscultation, OB-GYNs and other obstetric care providers and facilities should consider adopting protocols and training staff to use a hand-held Doppler device for low-risk women who desire such monitoring during labor.”⁴ • “Multiple nonpharmacologic and pharmacologic techniques can be used to help women cope with labor pain.”⁴ • “...use of the coping scale [rather than the 1-10 pain scale] can help OB-GYNS...tailor interventions to best meet the needs of each woman.”⁴ • “Women in spontaneously progressing labor may not require continuous infusion of IV fluids.”⁴ • Not an official position statement, but in this systematic review published in <i>Obstetrics and Gynecology</i>: “Women with low-risk singleton pregnancies who were allowed to eat freely during labor had a shorter duration of labor. A policy of less restrictive food intake during labor did not influence other obstetric or neonatal outcomes nor did it increase the incidence of vomiting. Operative delivery rates similar.”⁶
<p>5 Avoid giving birth on your back and follow your body’s urge to push.</p>	<ul style="list-style-type: none"> • “...each woman should be encouraged to use the technique [spontaneous versus Valsalva] that she prefers and is most effective for her.”⁴ <p>Although ACOG has supported the concept of delayed pushing (laboring-down) in the past, following the publication of a new study in JAMA⁷ in October of 2018, ACOG issued a practice advisory⁸ stating that, “it is reasonable to choose immediate over delayed pushing in nulliparous patients with neuraxial anesthesia.”</p>
<p>6 Keep your baby with you – it’s best for you, your baby, and breastfeeding.</p>	<p>“The Ten Steps [including early skin-to-skin care and rooming -in] should be integrated into maternity care to increase the likelihood that a woman will initiate and sustain breastfeeding...Skin-to-skin is feasible in the operating room and is associated with reduced need for formula supplementation.”⁹</p>

REFERENCES

1. Grobman, W.A., et al. (2018). Labor induction versus expectant management in low-risk nulliparous women. *New England Journal of Medicine*, 379(6), 513-523. PubMed abstract: www.ncbi.nlm.nih.gov/pubmed/30089070
2. American College of Obstetricians and Gynecologists. (2018). *Practice advisory: Clinical guidelines for integration of the findings of THE ARRIVE Trial: Labor induction versus expectant management in low-risk nulliparous women*. <https://bit.ly/2qltsnA>
3. American College of Obstetricians and Gynecologists and the Society for Maternal-Fetal Medicine. (2018). *Leaders in obstetric care respond to the published results of the ARRIVE Trial*. <https://bit.ly/2PyfwVB>
4. American College of Obstetricians and Gynecologists (ACOG). (Endorsed by ACNM & AWHONN). (2017). Committee Opinion #687: Approaches to limit intervention during labor and birth. *Obstetrics & Gynecology*, 129(2), 1-9. <https://bit.ly/2qIK6DI>
5. American College of Obstetricians and Gynecologists (ACOG) and the Society for Maternal-Fetal Medicine (SMFM). (2014, reaffirmed 2016). Obstetric care consensus: Safe prevention of the primary cesarean delivery. *Obstetrics & Gynecology*, 123(3), 693-711. <https://bit.ly/2yKdAQu>
6. Ciardulli, A., Saccone, G., Anastasio, H. & Berghella, V. (2017). Less-restrictive food intake during labor in low-risk singleton pregnancies. A systematic review and meta-analysis. *Obstetrics and Gynecology*, 129(3), 473-480. PubMed abstract: <https://bit.ly/2RsevMc>
7. Cahill, A.G., et al. (2018). Effect of immediate vs. delayed pushing on rates of spontaneous vaginal delivery among nulliparous women receiving neuraxial analgesia. A Randomized clinical trial. *JAMA*, 320(14), 1444-1454. PubMed abstract: <https://bit.ly/2OhEzYx>
8. American College of Obstetricians and Gynecologists (Endorsed by the American College of Nurse-Midwives and the Society for Maternal-Fetal Medicine). (2018). *Practice advisory: Immediate versus delayed pushing in nulliparous women receiving neuraxial analgesia*. <https://bit.ly/2yewmiC>
9. American College of Obstetricians and Gynecologists (ACOG). (2016). Committee opinion #658: Optimizing support for breastfeeding as part of obstetric practice. *Obstetrics and Gynecology*, 127(2), 1-7.